

TRYPTIC SOY AGAR W/ PENASE + 4 NEUTRALISANT (TRIPLE BAG)

INTENDED USE:

Tryptic Soy Agar (TSA, Casein Soya Bean Digest Agar) is a complex medium for the cultivation and isolation of fastidious bacteria, yeasts and moulds.

PRINCIPLE AND INTERPRETATION:

Tryptic Soy Agar (TSA, Casein Soya Bean Digest Agar) is a complex medium for the cultivation and isolation of fastidious bacteria, yeasts and moulds. The formulation of the basic medium is prepared according to Pharmacopoeial Harmonised Method and ISO Standards.

Tryptic Soy Agar with Penicillinase and neutralizers is used for Hygiene Monitoring (Environmental Monitoring) of air (active air samplers) on surfaces and personnel in the presence of penicillins.

The addition of the neutralizing agents TLHTh (Tween 80- Lecithin - Histidine - Sodium Thiosulphate) may inactivate a variety of disinfectants.

- * The combination of lecithin, polysorbate 80 and histidine neutralizes aldehydes and phenolic compounds.
- * The combination of lecithin and polysorbate 80 neutralizes the quaternary ammonium compounds.
- * The polysorbate 80 neutralizes hexachlorophene and mercurial derivatives.
- * Sodium thiosulphate neutralizes halogen compounds.
- * Lecithin neutralizes chlorhexidine.
- * Histidine neutralizes formaldehyde.

COMPOSITION:

Ingredients	Gr/Liter
Peptone from casein	15 gr
Soya peptone	5 gr
Sodium chloride	5 gr
Tween 80	5 gr
Lecithin	0,7 gr
Histidine	1 gr
Sodium Thiosulphate	0,5 gr
Penicillinase	10.000.000 IU
Agar	15 gr

***Formula adjusted, standardized to suit performance parameters

pH: 7,3 ± 0,2

PRECAUTIONS:

For professional use only. Do not use plates if they show evidence of microbial contamination, discoloration, drying, cracking or other signs of deterioration.

TEST PROCEDURE:

If the agar plate has been refrigerated, allow to warm to room temperature before inoculation. Mediums used in environmental control (active air, surface control, open petri dishes and personnel) are incubated for 48 hours at 30-35 ° C and 72 hours at 20-25 ° C.

QUALITY CONTROL:**1.Sterility Control:**

Incubation 3d at 30-35°C and 5d at 30-35°C: NO GROWTH

2.Physical/Chemical Control

pH: 7,3 ± 0,2

Appearance: Light amber

3.Microbiological Control: Incubation 72 hours at 20-25°C and 48 hours at 30-35°C

Microorganism	Inoculum (CFU)	Results
		Growth
Staphylococcus aureus ATCC 6538	10-100	Good
Escherichia coli ATCC 8739	10-100	Good
Pseudomonas aeruginosa ATCC 9027	10-100	Good
Candida albicans ATCC 10231	10-100	Good
Aspergillus brasiliensis ATCC 16404	10-100	Good
Bacillus subtilis ATCC 6633	10-100	Good
Staphylococcus epidermidis ATCC 12228	10-100	Good

STORAGE CONDITIONS AND SHELF LIFE:

Store the prepared medium at 2-12°C or 2-25°C. Use before expiry date on the label. Do not use beyond stated expiry date.

DISPOSAL:

Incubated prepared medium may contain active bacteria and micro-organisms. Do not open infected medium. Infected plate should be autoclaved, incinerated or opened and soaked in a chlorine-based disinfectant (liquid bleach) for 20 minutes prior to disposal.

PACKAGING:

Katalog Number: 02086

Packaging: Triple wrap/Gamma irradiated

Content: 10 plates/each package

REFERENCES:

- Abbott J. D. and Graham J. M. (1961) Mon. Bull. Min. Hlth Pub. Hlth Lab. Serv. 20. 51-58.
- Barrow G. I. and Ellis C. (1962) Mon. Bull. Min. Hlth Pub. Hlth Lab. Serv. 21. 141-147.
- Cooke G. T. and Daines C. F. (1964) Mon. Bull. Min. Hlth Publ. Hlth Lab. Serv. 23. 81-85.
- Gillies R. R. (1964) J. Hyg. Camb. 62. 1-9.
- Mitchell T. G. (1964) J. Appl. Bact. 27. 45-52.
- Barnes Ella M. and Shrimpton D. H. (1958) J. Appl. Bact. 2. 313-329.
- American Public Health Association (1978) Standard Methods for the Examination of Dairy Products. 14th Edn. APHA Inc. Washington DC.
- Anon. (1987) J. Food Microbiol. 5. 291-296.
- Lee K., Baron E.J., Summanen P. and Finegold S. (1990) J. Clin. Microbiol. 28. 1747-1750.
- Beumer R.R., te Giffel M.C. and Cox L.J. (1997) Lett. Appl. Microbiol. 24. 421-425.
- British Pharmacopoeia Volume II (2000)
- US Pharmacopoeia XXX, (2008)
- European Pharmacopoeia. 6.1 Edition (2008)
- Japanese Pharmacopoeia. 15th Edition. (2006)



Aseptik Steril



LOT Numarası



Referans Numarası



Negatif Kontrol



Pozitif Kontrol



Son Kullanma Tarihi



Sıcaklık Aralığı



Tek Kullanımlıktır



Ürün Adeti



Kullanım Kılavuzuna Bakınız



Üretici Firma